

# The routine immunisation schedule

from Autumn 2017

Age due	Diseases protected against	Vaccine given and trade name		Usual site <sup>1</sup>
Eight weeks old	Diphtheria, tetanus, pertussis (whooping cough), polio, <i>Haemophilus influenzae</i> type b (Hib) and hepatitis B	DTaP/IPV/Hib/HepB	Infanrix hexa	Thigh
	Pneumococcal (13 serotypes)	Pneumococcal conjugate vaccine (PCV)	Prevenar 13	Thigh
	Meningococcal group B (MenB)	MenB	Bexsero	Left thigh
	Rotavirus gastroenteritis	Rotavirus	Rotarix	By mouth
Twelve weeks old	Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B	DTaP/IPV/Hib/HepB	Infanrix hexa	Thigh
	Rotavirus	Rotavirus	Rotarix	By mouth
Sixteen weeks old	Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B	DTaP/IPV/Hib/HepB	Infanrix hexa	Thigh
	Pneumococcal (13 serotypes)	PCV	Prevenar 13	Thigh
	MenB	MenB	Bexsero	Left thigh
One year old *on or after the child's first birthday	Hib and MenC	Hib/MenC	Menitorix	Upper arm/thigh
	Pneumococcal	PCV	Prevenar 13	Upper arm/thigh
	Measles, mumps and rubella (German measles)	MMR <sup>3</sup>	MMR VaxPRO <sup>3</sup> or Priorix	Upper arm/thigh
	MenB	MenB booster	Bexsero	Left thigh
Two to eight years old <sup>2</sup> (including children in reception class and school years 1-4)	Influenza (each year from September)	Live attenuated influenza vaccine LAIV <sup>4</sup>	Fluenz Tetra <sup>3</sup>	Both nostrils
Three years four months old or soon after	Diphtheria, tetanus, pertussis and polio	DTaP/IPV	Infanrix IPV or Repevax	Upper arm
	Measles, mumps and rubella	MMR (check first dose given) <sup>3</sup>	MMR VaxPRO <sup>3</sup> or Priorix	Upper arm
Girls aged 12 to 13 years	Cervical cancer caused by human papillomavirus (HPV) types 16 and 18 (and genital warts caused by types 6 and 11)	HPV (two doses 6-24 months apart)	Gardasil	Upper arm
Fourteen years old (school year 9)	Tetanus, diphtheria and polio	Td/IPV (check MMR status)	Revaxis	Upper arm
	Meningococcal groups A, C, W and Y disease	MenACWY	Nimenrix or Menveo	Upper arm
65 years old	Pneumococcal (23 serotypes)	Pneumococcal polysaccharide vaccine (PPV)	Pneumococcal polysaccharide vaccine	Upper arm
65 years of age and older	Influenza (each year from September)	Inactivated influenza vaccine	Multiple	Upper arm
70 years old	Shingles	Shingles	Zostavax <sup>3</sup>	Upper arm <sup>5</sup>

<sup>1</sup> Where two or more injections are required at once, these should ideally be given in different limbs. Where this is not possible, injections in the same limb should be given 2.5cm apart. For more details see Chapters 4 and 11 in the Green Book. All injected vaccines are given intramuscularly unless otherwise stated.

<sup>2</sup> Age on 31 August 2017

<sup>3</sup> Contains porcine gelatine

<sup>4</sup> If LAIV (live attenuated influenza vaccine) is contraindicated and child is in a clinical risk group, use inactivated flu vaccine

<sup>5</sup> This can be administered subcutaneously but intramuscular is preferred.

All vaccines can be ordered from [www.immform.dh.gov.uk](http://www.immform.dh.gov.uk) free of charge except influenza for adults and Pneumococcal polysaccharide vaccine.

## Selective immunisation programmes

Target group	Age and schedule	Disease	Vaccines required
Babies born to hepatitis B infected mothers	At birth, four weeks and 12 months old <sup>1,2</sup>	Hepatitis B	Hepatitis B (Engerix B/ HBvaxPRO)
Infants in areas of the country with TB incidence $\geq$ 40/100,000	At birth	Tuberculosis	BCG
Infants with a parent or grandparent born in a high incidence country <sup>2</sup>	At birth	Tuberculosis	BCG
Pregnant women	During flu season At any stage of pregnancy	Influenza	Inactivated flu vaccine
Pregnant women	From 20 weeks gestation <sup>4</sup>	Pertussis	dTaP/IPV (Boostrix-IPV or Repevax)

1 Take blood for HBsAg at 12 months to exclude infection

2 In addition hexavalent vaccine (Infanrix hexa) is given at 8, 12 and 16 weeks

3 Where the annual incidence of TB is  $\geq$  40/100,000 – see <https://www.gov.uk/government/publications/tuberculosis-tb-by-country-rates-per-100000-people>

4 Can be given from 16 weeks but usually offered after the anomaly scan

## Additional vaccines for individuals with underlying medical conditions

Medical condition	Diseases protected against	Vaccines required <sup>1</sup>
Asplenia or splenic dysfunction (including due to sickle cell and coeliac disease)	Meningococcal groups A, B, C, W and Y Pneumococcal <i>Haemophilus influenzae</i> type b (Hib) Influenza	Hib/MenC MenACWY MenB PCV13 (up to two years of age) PPV (from two years of age) Annual flu vaccine
Cochlear implants	Pneumococcal	PCV13 (up to two years of age) PPV (from two years of age)
Chronic respiratory and heart conditions (such as severe asthma, chronic pulmonary disease, and heart failure)	Pneumococcal Influenza	PCV13 (up to two years of age) PPV (from two years of age) Annual flu vaccine
Chronic neurological conditions (such as Parkinson's or motor neurone disease, or learning disability)	Pneumococcal Influenza	PCV13 (up to two years of age) PPV (from two years of age) Annual flu vaccine
Diabetes	Pneumococcal Influenza	PCV13 (up to two years of age) PPV (from two years of age) Annual flu vaccine
Chronic kidney disease (CKD) (including haemodialysis)	Pneumococcal (stage 4 and 5 CKD) Influenza (stage 3, 4 and 5 CKD) Hepatitis B (stage 4 and 5 CKD)	PCV13 (up to two years of age) PPV (from two years of age) Annual flu vaccine Hepatitis B
Chronic liver conditions	Pneumococcal Influenza Hepatitis A Hepatitis B	PCV13 (up to two years of age) PPV (from two years of age) Annual flu vaccine Hepatitis A Hepatitis B
Haemophilia	Hepatitis A Hepatitis B	Hepatitis A Hepatitis B
Immunosuppression due to disease or treatment <sup>3</sup>	Pneumococcal Influenza	PCV13 (up to two years of age) <sup>2</sup> PPV (from two years of age) Annual flu vaccine
Complement disorders (including those receiving complement inhibitor therapy)	Meningococcal groups A, B, C, W and Y Pneumococcal <i>Haemophilus influenzae</i> type b (Hib) Influenza	Hib/MenC MenACWY MenB PCV13 (to any age) PPV (from two years of age) Annual flu vaccine

1 Check relevant chapter of green book for specific schedule

2 To any age in severe immunosuppression

3 Consider annual influenza vaccination for household members and those who care for people with these conditions